

Air Monitoring Summary Tables

The table below summarizes monitoring data collected using a portable wireless remote monitoring system. All times in Eastern Standard Time (EST).

From: 5/07/22 12:00 am **To:** 5/07/22 11:59 pm

Offsite Monitors

Instrument	Analyte	ATSDR MRL 14-day Avg Reached?	Concentration Range Detected ^a	24-hr Average ^a	7-day Average	ATSDR 14-day MRL
Catawba Headstart						
Acrulog PPB	H ₂ S	No	0 – 0 ppb	0.00 ppb	0.06 ppb	70 ppb
Treetops						
Acrulog PPB	H ₂ S	No	0 – 0 ppb	0.00 ppb	0.11 ppb	70 ppb
Liberty Hill						
Acrulog PPB	H ₂ S	No	0 – 0 ppb	0.00 ppb	0.07 ppb	70 ppb
Riverchase Estates						
Acrulog PPB	H ₂ S	No	0 – 4 ppb	0.10 ppb	0.28 ppb	70 ppb
Millstone Creek						
Acrulog PPB	H ₂ S	No	0 – 1 ppb	0.02 ppb	0.07 ppb	70 ppb

Onsite Fenceline Monitors


Instrument	Analyte	30-min AEGL Reached?	Concentration Range Detected ^a	24-hr Average ^a	7-day Average	30-min AEGL
Station 1						
TAPI Analyzer	H ₂ S	No	0 – 3 ppb	0.90 ppb	0.78 ppb	600 ppb
Station 2						
TAPI Analyzer	H ₂ S	No	1 – 8 ppb	2.23 ppb	1.51 ppb	600 ppb
Station 3						
TAPI Analyzer	H ₂ S	No	0 – 7 ppb	1.10 ppb	1.69 ppb	600 ppb


^a Based on 30-minute averages.

Notes:




- ATSDR MRL Agency for Toxic Substances and Disease Registry Minimal Risk Level (MRL)
- AEGL EPA Acute Exposure Guidelines Levels
- H₂S Hydrogen Sulfide
- TAPI Teledyne API H₂S Analyzer
- hr Hour
- min Minute
- ppb Parts per billion
- MRL Limit Limit defined as a 14-day average value.

Wind rose – Shows the direction the wind is coming from, the monitoring station being at the center of the rose.





Legend

-  Offsite Fixed Monitoring Locations
-  Onsite Fixed Monitoring Locations
-  New-Indy Catawba


Period H₂S Monitoring Hydrogen Sulfide Offsite Monitors

Below are graphs for offsite locations where hydrogen sulfide (H₂S) was detected during the current reporting period.


The five stand-alone H₂S monitoring stations correlate with five previous EPA's Viper monitoring system which includes areas to the north-northeast and south-southwest of the New-Indy Catawba Mill.

Winds came mostly from the southwest at 2 to 10 mph during the day and from the northeast at 4 to 15 mph during the night.


See wind rose diagram with aerial map figure for full wind data during this reporting period.




Treetops - 5/7/2022




Liberty Hill - 5/7/2022



Riverchase Estates - 5/7/2022



Millstone Creek - 5/7/2022





Period H₂S Monitoring Hydrogen Sulfide Onsite Monitors

Below are graphs for onsite locations during the current reporting period.

Depending on wind direction, the H₂S measured at the onsite fence line locations may not exit the mill property at reported concentrations. Wind directions from offsite locations, blowing onto mill property, will disperse ambient concentrations to lower levels prior to exiting the plant site.

Winds came mostly from the southwest at 2 to 10 mph during the day and from the northeast at 4 to 15 mph during the night.

See wind rose diagram with aerial map figure for full wind data during this reporting period.



Station3 - 5/7/2022

